Carbohydrate Research, 252 (1994) C9–C11
Elsevier Science B.V., Amsterdam

CONTENTS

An NMR spectroscopic and conformational study of 12 pseudo-disaccharides (D-glucopyranosyl-5a-carba-D- and -L-glucopyranoses)
J. Duus, K. Bock (Copenhagen Valby, Denmark) and S. Ogawa (Yokohama, Japan) 1

The crystal structure of the inclusion complex of cyclomaltoheptaose (β-cyclodextrin) with 4-*tert*-butylbenzoic acid
A. Rontoyianni, I.M. Mavridis, E. Hadjoudis (Athens, Greece) and A.J.M. Duisenberg (Utrecht, Netherlands) 19

Effect of heat–moisture treatment on the structure and physicochemical properties of cereal, legume, and tuber starches
R. Hoover and T. Vasanthan (St. John's, NF, Canada) 33

Towards the selective acylation of secondary hydroxyl groups of carbohydrates using oxime esters in an enzyme-catalyzed process
R. Pulido and V. Gotor (Oviedo, Spain) 55

Chemical synthesis and NMR spectra of a protected branched-tetrasaccharide thioglycoside, a useful intermediate for the synthesis of branched oligosaccharides
M.S. Motawia, C.E. Olsen, B.L. Møller and J. Marcussen (Copenhagen, Denmark) 69

2-Acetamido-2-deoxyaldonolactones from sugar formazans
V. Zsoldos-Mády, I. Pintér, A. Neszmélyi, A. Messmer and A. Perczel (Budapest, Hungary) 85

Synthesis, crystal structure, and conformation of methyl 6-deoxy-2,3-*O*-isopropylidene-α-D-*manno*-heptofuranoside
J.W. Krajewski, P. Gluziński, Z. Pakulski, A. Zamojski (Warszawa, Poland), A. Mishnev and A. Kemme (Riga, Latvia) 97

Synthesis of α and β anomers of UDP-*N*-acetylmuramic acid
D. Blanot, G. Auger, D. Liger and J. Van Heijenoort (Orsay, France) 107

Synthesis of the globotetraose tetrasaccharide and terminal tri- and di-saccharide fragments
U. Nilsson, A.K. Ray and G. Magnusson (Lund, Sweden) 117

Synthesis of the Forssman pentasaccharide and terminal tetra-, tri-, and di-saccharide fragments
U. Nilsson, A.K. Ray and G. Magnusson (Lund, Sweden) 137

Solvent-free benzylation of polyols by phase-transfer catalysis or supported reagent methods
D. Dubreuil, J. Cleophax (Gif-sur-Yvette, France) and A. Loupy (Orsay, France) 149

Synthesis of di- and tri-saccharides with intramolecular NH-glycosidic linkages: molecules with flexible and rigid glycosidic bonds for conformational studies
J. Goddat, A.A. Grey, M. Hricovíni, J. Grushcow, J.P. Carver and R.N. Shah (Toronto, ON, Canada) 159

Carrageenan from the tetrasporic stage of *Gigartina decipiens* (Gigartinaceae, Rhodophyta)
R. Falshaw and R.H. Furneaux (Lower Hutt, New Zealand) 171

Structural roles of calcium ions and side chains in welan: an X-ray study
R. Chandrasekaran, A. Radha and E.J. Lee (West Lafayette, IN, USA) 183

Native ordered structure of welan polysaccharide: conformational transitions and gel formation in aqueous dimethyl sulphoxide
M.W.N. Hember, R.K. Richardson and E.R. Morris (Bedford, United Kingdom) 209

Structural elucidation of the specific capsular polysaccharide of *Rhodococcus equi* serotype 7
H. Masoud and J.C. Richards (Ottawa, ON, Canada) 223

Isolation and structural characterization of β-D-glucosyluronic acid and 4-*O*-methyl β-D-glucosyluronic acid-containing oligosaccharides from the cell-wall pectic polysaccharide, rhamnogalacturonan I
J. An, M.A. O'Neill, P. Albersheim and A.G. Darvill (Athens, GA, USA) 235

Isolation and characterisation of cell wall polymers from olive pulp (*Olea europaea* L.)
M.A. Coimbra, (Aveiro, Portugal), K.W. Waldron and R.R. Selvendran (Norwich, United Kingdom) 245

Notes

Determination of the absolute configuration of the glycerol component in poly(glycosylglycerolphosphates) by GLC–MS
L.M. Beynon and J.C. Richards (Ottawa, ON, Canada) 263

X-ray structure of methyl 1,6-di-*O*-(4-bromobenzoyl)-3,4-di-*O*-(4-methoxycinnamoyl)-β-D-fructofuranoside
N. Ikemoto (New York, NY, USA) and G.K. Schulte (New Haven, CT, USA) 269

Synthesis of D-erythrofuranosyl C-nucleosides of imidazole from 4(5)-(D-*arabino*-tetritol-1-yl)imidazole
D. Deredas and A. Frankowski (Łódź, Poland) 275

A facile total synthesis of ganglioside GD2
H. Ishida (Gifu, Japan), Y. Ohta, Y. Tsukada (Uji, Japan), Y. Isogai, H. Ishida, M. Kiso and A. Hasegawa (Gifu, Japan) 283

Oxidation of sugar hydroxy groups with triphenylphosphine–*N*-chlorosuccinimide–dimethyl sulfoxide
G. Hodosi (Budapest, Hungary) 291

A convenient laboratory procedure for the preparation of cortalcerone, a fungal antibiotic β-pyrone
J. Gabriel, J. Volc, E. Kubátová, Z. Palková and M. Pospíšek (Prague, Czech Republic) .. 297

L-Dianose, a new monosaccharide from *Dianthus chinensis* L.
H.-y. Li (Chengdu, P.R. China), K. Koike, T. Ohmoto (Chiba, Japan), Y.-s. Li and G.-f. Guo (Liaoyuan, P.R. China) 303

Structure of the capsular polysaccharide and the O-side-chain of the lipopolysaccharide from *Acetobacter methanolicus* MB 58 (IMET B346)
H.D.G.U. Mamat, I. Kühn (Leipzig, Germany), A.S. Shashkov and Y.A. Knirel (Moscow, Russian Federation) 309

Structural and molecular properties of the arabinogalactan isolated from Mongolian larchwood (*Larix dahurica* L.)
P. Odonmažig (Ulan-Bator, Mongolia), A. Ebringerová, E. Machová and J. Alföldi (Bratislava, Slovakia) 317

Differing rates of galactosylation of homologous spacer-modified disaccharides are an indication for two separate GlcNAc binding-sites in galactosyltransferase
S.-C. Ats, J. Lehmann and S. Petry (Freiburg, Germany) 325

Book review C1

Author index C3

Subject index C5

Contents C9